

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

SWA4338P0080US (WP/1801)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

10/089491

INTERNATIONAL APPLICATION NO.
PCT/CA 00/01118INTERNATIONAL FILING DATE
28/09/2000PRIORITY DATE CLAIMED
28 September, 2000 (22.09.00)

TITLE OF INVENTION

HUMAN FMO3 GENE MUTATIONS AND POLYMORPHISMS

APPLICANT(S) FOR DO/EO/US

Eileen P. Treacy; Beverly R. Ackerman

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☐ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☐ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☒ Certificate of Mailing by Express Mail Label No. EM306837149US
23. ☒ Other items or information:

Paper Copy of Sequence Listing

Copy of International Publication Number WO 01/23603 A2

Copy of Form PCT/IB/308

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.101) 10/089491		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER SWA4338P0080US	
---	--	-------------------------------	--	---	--

24. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) : <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY <div style="border: 1px solid black; height: 100px; width: 100%;"></div>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).				\$890.00 \$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	18 - 20 =	0	x \$18.00	\$0.00	
Independent claims	5 - 3 =	2	x \$84.00	\$168.00	
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>				\$0.00	
TOTAL OF ABOVE CALCULATIONS =				\$1,058.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$0.00	
SUBTOTAL =				\$1,058.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				\$0.00	
TOTAL NATIONAL FEE =				\$1,058.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00	
TOTAL FEES ENCLOSED =				\$1,058.00	
				Amount to be: refunded	\$
				charged	\$

a. ☒ A check in the amount of **\$1,058.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. **23-0785**. A duplicate copy of this sheet is enclosed.


d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Martin L. Katz
 Wood, Phillips, Katz, Clark & Mortimer
 Citicorp Center, Suite 3800
 500 W. Madison St.
 Chicago, IL 60661-2511

Telephone: (312) 876-1800
 Facsimile: (312) 876-2020


 SIGNATURE

Martin L. Katz
 NAME

25,011
 REGISTRATION NUMBER

March 28, 2002
 DATE

Docket No. SWA4338P0080US

Corinne Byk

PATENT
Docket No. SWA4338P0080US

REMARKS

Applicants have amended claim 5 to delete the multiple dependency of claim 5, thereby eliminating the multiple dependency of all claims dependent on claim 5.

Thus, the total number of claims is 18, with 3 independent claims with a fee of \$529 for an applicant with small entity status. No surcharge for multiple dependent or excess claims should now be applicable.


Should there be any additional fees required, the Commissioner is hereby authorized to charge Deposit Account No. 23-0785.

Respectfully submitted,

WOOD, PHILLIPS, KATZ, CLARK & MORTIMER

Date: December 16, 2002

By



Martin L. Katz
Reg. No. 25,011

500 West Madison Street
Suite 3800
Chicago, IL 60661
(312) 876-1800

PATENT
Docket No. SWA4338P0080US

Marked-Up Version of Claims

5. (amended) A method according to [anyone of] claim[s] 1 [to 4], wherein said at least one of said mutation and said polymorphic variant inactivates partially or totally the activity of said FMO enzyme.

HUMAN FMO3 GENE MUTATIONS AND POLYMORPHISMS,
AND USES THEREOF

BACKGROUND OF THE INVENTION

5 (a) Field of the Invention

The invention relates to human FMO3 gene polymorphisms and mutations, and more particularly to uses thereof in pharmacogenomics and the diagnosis of inborn errors of metabolism such as trimethylaminuria and predisposition to hypertension.

10 (b) Description of Prior Art

Inter-individual variations resulting in population-wide differences in the metabolism of foreign compounds or xenobiotics may contribute to the susceptibility of humans to adverse chemicals or drug reactions and disease states. Most incidences of genetic variation can be accounted for by more prevalent alleles such as alleles with a frequency of greater than 1% in the general population, called polymorphisms or common variants. It is probable that common variants may contribute in a significant fashion to disease susceptibility. Pharmacogenomics allows for the identification of genetic variation in drug-metabolizing enzymes and the identification of individuals who will benefit most or least from a given medication.

Detoxification of xenobiotics including drugs, food additives and environmental chemicals is mediated by Phase I (oxidative) and Phase II (conjugative) reactions. The microsomal cytochrome P450 (CYP) family of monooxygenases is highly polymorphic in humans. CYP3A4 is largely responsible for hepatic drug metabolism. However, CYP2D6 metabolizes a variety of widely prescribed drugs, and individuals with impaired CYP2D6 activity show a number of drug interactions

resulting from aberrant metabolism due to inactivating polymorphisms.

Human flavin-containing monooxygenases (FMO) (EC 1.14.13.8) are microsomal NADPH-dependent flavoprotein enzymes that catalyze the oxygenation of nucleophilic nitrogen-, sulfur-, phosphorus- and other heteroatom-containing chemicals, drugs and pesticides. The FMOs belong to the FSSP flavocytochrome c sulfide dehydrogenase subfamily of flavoenzymes, NAD(P)H-dependent monooxygenases and reductases. FMOs are membrane-bound proteins that have been detected in all secretory cell types that have been examined. Human FMOs are 532-558 amino acids in length, with specific amino acids highly conserved in all species, particularly residues 4-32 and 186-213, which contain the FAD and NADPH-binding domains, respectively (Cashman J (1995) Chem Res Toxicol 8:166-181). The amino acid sequence identity between human isoforms is at a minimum of 52%. FMO enzymes have a broad substrate specificity compared to other mammalian monooxygenases, with at least 1,000 known substrates for these enzymes. They represent a family of five monooxygenases in mammals designated FMO, forms 1-5.

Differences in drug metabolism among animal species have been recognized for more than 50 years. Human FMO drug metabolizing enzymes are thought to have evolved as a multi-gene family from an original monooxygenase such as that cloned and characterized from yeast (*Saccharomyces cerevisiae*) which has high amino acid sequence conservation with mammalian FMO.

Of the five FMO monooxygenase enzymes, FMO3 constitutes the prominent form that converts nucleophilic heteroatom-containing chemicals and endogenous materials to polar, readily excreted oxygenated metabolites, and this facilitates their

elimination. *FMO3* constitutes the major adult human hepatic isoform. Biogenic amines including primary amines, such as tyramine and phenylethylamine, and tertiary amines are good substrates for human *FMO3* (Cashman J (1995) Chem Res Toxicol 8:166-181). Tertiary amine substrates include trimethylamine (TMA), antihistamines, and (S)-nicotine. TMA and tyramine are N-containing substrates of the enzyme. Commonly used drugs such as tricyclic antipsychotics, cimetidine, ranitidine, albendazole and verapamil are also oxygenated by human *FMO3* (Cashman J (1995) Chem Res Toxicol 8:166-181, Ziegler, 1990).

The odorous, dietary-derived tertiary amine trimethylamine (TMA) is N-oxygenated by human *FMO3* to the non-odorous trimethylamine N-oxide (TMANO). TMANO is excreted in a detoxication and deodoration process. In normal humans, over 95% of TMA is metabolized to TMANO and is excreted in the urine at concentrations of less than 18 μmol of TMA/ mmol creatinine under normal dietary conditions. In a small number of humans, TMA is not efficiently metabolized to TMANO, and those individuals suffer from trimethylaminuria or fish-like odor syndrome, due to the presence of relatively large amounts of this odorous volatile amine in bodily excretions, including urine, sweat and breath.

Trimethylaminuria (TMAuria), first described in 1970, is an autosomal recessive inborn error of metabolism, which results in a partial or total inability to oxidize TMA to TMANO. (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998) and a severe body odor reminiscent of rotten fish emanating in sweat, breath and urine with associated psychosocial disorders. This condition, previously thought to be rare, is now being increasingly detected in severe and

milder presentations. TMAuria occurs in Australians and has a predicted incidence of ~1/40,000 in North Americans.

Current treatment strategies include dietary
5 restriction of TMA precursors (choline, lecithin and carnitine), the intermittent use of antibiotics such as metronidazole, and other supportive measures.

Treatment options for this condition are limited and not well studied. Current strategies
10 include limitation of choline and TMA precursors. Dietary restrictions are insufficient for a number of individuals (related to genotype). Also some individuals over restrict choline which may also cause side-effects.

15 A number of individuals are consuming severely protein-restricted diets. Choline is an essential amine required for synthesis of phosphatidylcholine, sphingomyelin, acetylcholine and betaine (essential for remethylation of homocysteine) and for
20 myelination. Choline deficiency causes memory impairment, liver and kidney dysfunction and cell death by apoptosis.

In 1998, choline was classified as an essential human nutrient by the Food and Nutrition Board of the
25 Institute of Medicine of the National Academy of Sciences. Choline intake requires careful assessment in children and expectant mothers. There is as yet no studies of choline requirements in individuals with TMAuria on and off diet, and the precise choline
30 limitation required for satisfactory decreases of TMA levels and relief of symptoms has never been studied. Choline and derivatives such as phosphocholine are present in many food stuffs including milk and infant formulas in a substantial amount (25). Precise
35 definition of choline requirements in individuals with

TMAuria would be useful to determine a choline-restricted formula.

5 The structure for the *FMO3* gene has been characterized and maps to chromosome 1q23-25. The *FMO3* gene is approximately 28 kb in length, with 1 noncoding and 8 coding exons (Gene Bank Accession Number AL021026).

10 Although substrates are generally detoxicated by *FMO*-mediated metabolism, the *FMO3* enzyme has been implicated in the bioactivation of a number of xenobiotics (Cashman J (1995) Chem Res Toxicol 8:166-181). Thus, inactivating variants or polymorphisms of human *FMO3* may contribute to the pathophysiology of diseases and adverse reactions or exaggerated clinical
15 responses to specific medications.

There exist genotyping assays which predict altered drug metabolism for the parallel phase I drug metabolizing enzyme system cytochrome P450.

20 There are at present no clinical tests available for pharmacogenomics with respect to *FMO3* and no markers for altered predisposition to handling of *FMO3* substrates.

It would therefore be highly desirable to be provided with a diagnostic test for trimethylaminuria.

25 It would also be highly desirable to be provided with identified pharmacogenomic polymorphisms of *FMO3*. The detection of same would allow the determination of differential responses to medications and environmental toxins related to diseases such as
30 those involving a complex pathophysiology. A large scale screening of the population with the detection of same would be advantageous to the pharmaceutical industry. The upregulation or downregulation of such polymorphic genotypes may also be made possible.

SUMMARY OF THE INVENTION

One aim of the present invention is to provide a diagnostic test for trimethylaminuria (TMAuria).

Another aim of the present invention is to provide identified polymorphisms of *FMO3*. The identified polymorphisms may allow the determination of differential responses to medications and environmental toxins causing diseases.

Mutations of the *FMO3* gene have been determined to cause trimethylaminuria.

Several single nucleotide polymorphisms (SNPs) of the *FMO3* gene have been identified. Rare mutations were discovered and identified use of the mutations and polymorphisms to determine their consequences for detoxication of drugs, chemicals and endogenous materials is also provided.

The distribution of the three polymorphisms V257M, E158K and E308G was studied in different combinations in Quebec individuals (Genetic Signatures). *In vitro* expression analysis in an *E. coli* system indicated that the codon 158 and 257 polymorphisms are pharmacogenomic, prevalent polymorphisms. Preliminary clinical evidence is provided to the effect that the E308G polymorphism, particularly when in *cis* with the variant codon 158 allele, alters oxidation of trimethylamine, likely applicable to other *FMO3* substrates. Analysis of cDNA expressed *FMO3*-maltose-binding fusion proteins for two of those polymorphisms demonstrated altered N-oxygenation for 10-(N'-N-dimethylaminopentoyl)-2-(trifluoromethyl) phenothiazine, indicating that these are significant pharmacogenomic polymorphisms.

Diagnostic tests for the mutations and polymorphisms are provided. The study of the distribution and population frequency of these three

polymorphisms (pharmacogenomics) may likely predict adverse or altered response to typical *FMO3* substrates and common medications and indicate predispositions to complex disorders associated with disordered
5 xenobiotic and biogenic amine metabolism. Expression analysis for the third polymorphism may be effected.

Loss of N-oxygenation and decreased clearance of amine-containing chemicals or drugs may contribute to adverse drug reactions, interactions with
10 catecholamine or exaggerated clinical response to specific medications.

The detection of *FMO3* polymorphisms may be used to determine differential responses to medications and environmental toxins applicable to complex disease
15 pathophysiology, and to determine high throughput screening for polymorphism detection (DNA chip technology) for large scale population, which may be advantageous to the pharmaceutical industry. Polymorphic *FMO3* genotypes may also be upregulated or
20 downregulated.

In accordance with the present invention, there is provided a method for detecting an altered metabolism of a substrate of a flavin-containing monooxygenase (*FMO*) enzyme or an isoform thereof in an
25 individual. The method comprises detecting at least one of a mutation and a polymorphic variant of a gene encoding the *FMO* enzyme or isoform thereof in a sample from the individual, whereby the at least one of the mutation and the polymorphic variant is indicative of
30 an altered metabolism for the substrate.

In accordance with the present invention, there is also provided a method for detecting a susceptibility of an individual to a substrate of the *FMO* enzyme or isoform thereof in an individual. The
35 method comprises detecting at least one of a mutation

and a polymorphic variant of a gene encoding the FMO enzyme or isoform thereof in a sample from the individual, whereby the at least one of the mutation and the polymorphic variant is indicative of a susceptibility to the substrate.

5 In accordance with the present invention, there is further provided a method for detecting a predisposition of an individual to a disorder associated with an (adverse) exposure to a heteroatom-containing chemical compound, an intermediate or a
10 metabolite thereof associated with carcinogenesis or having a toxic, pro-carcinogenic or carcinogenic potential. The method comprises detecting at least one of a polymorphic variant and a mutation of a gene
15 encoding the FMO enzyme or an isoform thereof in a sample from the individual, whereby the at least one of the polymorphic variant and the mutation is indicative of exposure to the chemical compound, the intermediate or the metabolite thereof.

20 In accordance with the present invention, there is further provided a method for detecting a predisposition to hypertension in a patient, the method comprising detecting at least one of a mutation and a polymorphic variant of a gene encoding the FMO
25 enzyme in a sample from the patient, whereby at least one of the mutation and the polymorphic variant is indicative of a predisposition to hypertension.

In accordance with the present invention, there is further provided a method for the treatment of an
30 individual having a disorder associated with an altered activity of the FMO enzyme or an isoform thereof. The method comprises supplementing the individual with riboflavin to increase the altered activity of the FMO enzyme or the isoform thereof.

The mutation or polymorphic variant may inactivate partially or totally the activity of the FMO enzyme.

5 The isoform of the enzyme may consist of form 3 (FMO3).

The polymorphic variant may comprise a polymorphic variant from the group consisting of E158K, V257M and E308G, and the mutation may comprise a mutation from the group consisting of P153L, E305X, 10 M66I, E314X, R492W, A52T and R387L.

The altered metabolism may be associated with an idiosyncratic reaction to the substrate.

The altered metabolism may be associated with a disorder such as a cancer. The disorder may also be 15 TMAuria.

The substrate may consist of a xenobiotic or an endogenous material relative to the individual. The xenobiotic may be a drug, a food additive, a pesticide, a plant toxin, an organic chemical compound 20 or an aromatic amine. The substrate may be a biogenic amine contained in the individual's diet. The biogenic amine may consist of a tertiary amine, such as trimethylamine (TMA), tyramine or catecholamine.

For the purpose of the present invention, the 25 following abbreviations are defined below.

"FMO" is intended to mean human flavin-containing monooxygenase;

"FMO3" is intended to mean human flavin-containing monooxygenase form 3;

30 "FMO1" is intended to mean human flavin-containing monooxygenase form 1;

"TMA" is intended to mean trimethylamine;

"TMANO" is intended to mean trimethylamine N-oxide;

"5-APT" is intended to mean 10-(5-Aminopentyl)- 35 2(trifluoromethyl)phenothiazine;

"5-DPT" is intended to mean 10-(N,N-Dimethylaminopentyl)-2-(trifluoromethyl)phenothiazine;
MBP: maltose binding protein;

5 "FMO3 MBP" is intended to mean human flavin-containing monooxygenase form 3 maltose binding protein;

"SDS-PAGE" is intended to mean sodium dodecyl sulfate polyacrylamide gel electrophoresis;

"TCA" is intended to mean trichloroacetic acid; and

"TMAuria" is intended to mean trimethylaminuria.

10 A "polymorphism" is intended to mean a nucleotide substitution of a gene that occurs at a frequency greater than 1% in individuals in the general population.

"Severe" TMAuria is intended to mean a
15 reduction of TMA oxidation below 50% of normal oxidation.

A "genetic signature" with respect to a substitution is intended to mean a combination of a polymorphism or a mutation identified in an individual
20 or haplotype for the substitution of interest.

The term "patient" is intended to mean a mammal preferably human.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Fig. 1 illustrates in a diagram the structural organization of FMO3 cDNA.

DETAILED DESCRIPTION OF THE INVENTION

Two mutations of the FMO3 gene, P153L and E305X
30 (OMIM accession numbers 136132.001 and 136132.004) account for greater than 90% of cases of TMAuria in Australians. Six other mutations (Fig. 1) and three intragenic polymorphisms (E158K, V257M and E308G) which appear to affect TMA and other FMO3 substrate
35 oxidation are also reported. In a North-American

cohort (including 4 Canadians) 100% ascertainment of mutant alleles was reported and in some probands, the second mutant allele comprises two of these polymorphisms in *cis*, (K158-G308) (Akerman BR et al. Am J Hum Genet (1997) Supplement to vol 61 (no. 4). Abs. no.281) (Table 1). Of the three polymorphisms identified in Quebec (see Table 2), expression of mutant cDNAs for the codon 158 and 257 polymorphisms showed altered N-oxygenation for 5-DPT (a phenothiazine substrate), tyramine and TMA, suggesting that these polymorphisms are significant prevalent pharmacogenomic polymorphisms.

A cohort of individuals was ascertained in North America with severe TMAuria.

Of 70 referrals for a suspected diagnosis of TMAuria, TMAuria was confirmed in 20 individuals with reduced TMANO/TMA ratios. (Normal values are: TMANO/TMA relative % ratios > 98:2). Among these, some had symptoms of hypertension and migraine with abnormal dopamine metabolism, suggesting that mutations of the *FMO3* gene may result in abnormal catecholamine metabolism.

Four new *FMO3* mutations were detected in this cohort: two missense (A52T and R387L), and one nonsense (E314X) (Akerman BR et al. Am J Hum Genet (1997) Supplement to vol 61 (no. 4). Abs. no.281). The fourth allele is apparently composed of two relatively common polymorphisms (K158-G308) found in the general population.

A number of prevalent nucleotide polymorphisms of the human *FMO3* gene were identified. Polymorphisms that are prevalent in French Canadian and Australian populations were studied. The cDNA-expression analysis for two of these prevalent human *FMO3* polymorphisms showed altered N-oxygenase activities, indicating that

these are significant pharmacogenomic polymorphisms.

Two prevalent polymorphisms of this gene (E158K and V257M) modulate the activity of human *FMO3*. These polymorphisms are widely distributed in Canadian and Australian Caucasian populations. *In vitro* analysis of wild-type and variant human *FMO3* proteins expressed from the cDNA for these two naturally-occurring polymorphisms showed differences in substrate affinities for nitrogen-containing substrates. Thus, for polymorphic forms of human *FMO3*, lower k_{cat}/K_m values for N-oxygenation of 10-(N,N-dimethylaminopentyl)-2-(trifluoromethyl)phenothiazine, trimethylamine and tyramine were observed. The results imply that prevalent polymorphisms of the human *FMO3* gene likely represent low penetrance predispositions to diseases associated with adverse environmental exposures to heteroatom-containing chemicals, drugs and endogenous amines.

Analysis of polymorphisms may be performed by PCR testing in Quebec residents with application to metabolism of *FMO3* substrates (medications) in humans that are heterozygous or homozygous for polymorphisms. The genotyping methods are described herein. The genotypes observed in Quebec (Genetic Signatures) and methods thereof are also described. Deficient human *FMO3* N-oxygenase activity has implications for the abnormal metabolism of many other endogenous dietary and medicinal amines.

We previously reported that the enzyme encoded by the human *FMO3* Glu158 polymorphic allele is more active for the tertiary amine substrates TMA and 5-DPT than the enzyme encoded by the less prevalent human *FMO3* Lys158 allele (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998).

Two new other polymorphisms are provided herein. The presence of these prevalent *FMO3* polymorphisms implies that some individuals may be more susceptible to the effects of environmental chemicals and to idiosyncratic drug reactions. Thus, if a drug or chemical is dependent on human *FMO3* for detoxication and if an individual possesses a impaired polymorphic form of *FMO3* then the drug or chemical may produce an exaggerated clinical response and this could lead to adverse reactions.

Human *FMO1* is not functionally active in adult human liver (Cashman J (1995) Chem Res Toxicol 8:166-181). Based on the kinetic parameters observed, human *FMO1* does not appear to make a significant contribution to the metabolism of TMA. From the present results it is likely that human *FMO1* does not 'rescue' an individual that is deficient in *FMO3* from suffering the consequences of trimethylaminuria. Human *FMO1* does not significantly N-oxygenate primary amines such as tyramine or 5-APT but can participate in the N-oxygenation of chemicals or drugs containing the tertiary amine functionality.

From the *in vitro* data, the codon 257 polymorphism appears to show substantial differences in kinetics for the biogenic amine substrate, tyramine and the dietary amine, TMA. This may have clinical consequences.

The maintenance of the prevalent codon 158 polymorphism in the Caucasian populations examined may not result solely from genetic drift, but perhaps as a consequence of molecular drive, whereby particular polymorphisms with selective advantages persist, for example, to combat exposure to plant toxins in particular geographic regions. The distribution of the two codon 158 alleles is almost in equilibrium in

these populations, suggesting that this may be an older polymorphism that is now balanced. Human *FMO3* V257M is of lower prevalence and may represent a founder effect.

5 Variation in human drug metabolism by genetic polymorphisms may increase the risk for acquiring exposure-related disease, including cancer, with important public health consequences. Generally, oxidative metabolism of heteroatom-containing
10 compounds by CYP-dependent processes leads to production of chemical intermediates with increased potential for toxic or carcinogenic properties. Although human *FMOs* generally convert lipophilic heteroatom-containing compounds to polar, readily
15 excreted oxygenated metabolites that possess decreased toxic potential, they may also catalyze the N-oxygenation of a wide array of xenobiotics such as plant toxins, organic chemicals and aromatic amines associated with carcinogenesis (Cashman J (1995) *Chem*
20 *Res Toxicol* 8:166-181).

Consistent with the fact that human drug-metabolizing enzymes have endogenous substrates and are prevalent not as neutral balanced polymorphisms but for their selective advantages, it was previously
25 shown that human *FMO3* metabolizes biogenic amines such as tyramine and phenethylamine resulting in great stereoselectivity of their oxime metabolites. Formation of oxime metabolites generally terminates the pharmacological activity of the parent amine (Lin, J et al. (1997) *Chem. Res. Toxicol.* 10: 842-852). It
30 is shown herein that the methionine variant at codon 257 of human *FMO3* shows decreased N-oxygenation for the substrate tyramine. Tyramine is an indirectly-acting sympathomimetic that exerts its pressor effect
35 through amine uptake into the sympathetic nervous

system with release of norepinephrine from synaptic vesicles. It is thus possible that human *FMO3* polymorphisms affecting tyramine or other biogenic amine metabolism may predispose humans to variable tolerance to tyramine or other biogenic amine-containing foods and associated symptoms.

It is shown that human *FMO3* null alleles are rare and cause a severe phenotype with lack of oxygenation of human *FMO3* substrates such as TMA (Cashman JR et al. Chem Res Toxicol 10:837-841, 1997; Treacy EP et al. Hum Molec Genet 7: 839-845, 1998). On the basis of the in vitro data described herein, the human *FMO3* polymorphisms have been shown to have milder effect on N-oxygenation activity. The polymorphisms are, however, of higher frequency, and thus they are more likely to show specific associations with complex diseases in populations.

The present invention will be more readily understood by referring to the following examples which are given to illustrate the invention rather than to limit its scope.

EXAMPLE I

TMAuria is caused by mutations of the *FMO3* gene in a North American Cohort

Urine detection

Early morning first void samples were collected and acidified with N HCl to pH 2 upon collection from subjects, and TMA and TMANO were measured by fast atom bombardment mass spectrometry using ¹⁶N-labeled TMA and TMANO as internal standards, a highly sensitive assay.

Values were expressed as mmole TMA or TMANO per mole creatinine. Control ratios for 20 adult urine collected from volunteers were established as TMANO:TMA percentage ratios >98:2%.

Mutation detection

DNA was extracted by standard methods from whole blood collected in EDTA from probands with TMAuria. The samples were then screened by PCR and restriction digestion for amino acid polymorphisms previously identified (E158K, V257M and E308G) and for mutations detected in the Australian TMAuria cohort (M66I, P153L, E305X, R492W).

Table 1 lists conditions used for the restriction diagnostic tests performed and the relevant primer sequences.

Table 1
Diagnostic PCR Reactions
for human HFMO3 Variants

Allele	Exon	Primers (sense-antisense; 5' to 3')	SEQ ID No:	Restriction enzyme	Expected normal (bp)	Pattern mutant (bp)
A52T	3	2038: GACCTGATCAGTATACTCATTTA 2025: AATGGGAAGTCTGGGAAACACTT	1	<i>NheI</i>	79	147
M66I	3	As above	2	<i>MseI</i>	147	122
P153	4	2055: TTGTTCCGGACATCATGTGTAGC Gen2: TCCCTGCTGTGGAAGCATTT	3	<i>AluI</i>	263	263
K158E	4	As above	4		66	45
				<i>HinfI</i>	253	217
					76	76
V257M	6	C1: TTCCAGAAGTGGCTCCTGGG 2005: GCTTGAATCTTGCAATCATCTGC	5	<i>Hsp92II</i>	97	36
			6		60	72
E305X	7	2019: CCTTATCAATTTATATATGGACC 2016: GGACCTTGTAACATAGGATTATTG	7	<i>EcoRI</i>	365	25
E308G	7	As above	8		164	529
				<i>BsaI</i>	362	529
E314X	7	As above			167	
				<i>AclI</i>	409	532
R387L	7	As above			122	
				<i>MseI</i>	112	102
R492W	9	2049: GAAATGCCATACTGACCCCATGG 2050: TAGCAAAGCCCCTGTCTGGGTAT	9	<i>BstXI</i>	102	88
			10		211	24
						186
						25

To confirm the three newly identified mutations, restriction enzyme analysis was performed on three independent PCR products from the probands. For the A52T and R387L alleles, the mutations ablated naturally-occurring restriction sites, while the E314X variant created a new restriction site, as shown in

Table 1. The mutation assays were subsequently used to determine the frequency of mutations in control samples from the province of Quebec, Canada.

Results

5 Of 28 individuals referred for investigation of
TMAuria, we identified 10 who had a marked decrease in
TMA oxidation, with less than 50% of the total TMA in
the N-oxide form, as shown in (Table 2). Control
10 ratios for these compounds were defined as being
greater than 98:2% in our laboratory. Further analysis
was possible for 10 of the remaining undiagnosed
consultants by gas chromatography-mass spectrometry;
no abnormalities were detected. In addition, testing
15 of these 10 urine samples by NMR spectroscopy ruled
out dimethylglycinuria, a recently described entity
which also presents with a fish-like odor and mild
TMAuria.

 Four of the parents of our defined TMAuria
probands (obligate heterozygotes) were also tested for
20 TMANO:TMA ratios. For one heterozygote carrying the
E314X mutation the TMANO:TMA ratio was noted to be
95:5. For the other heterozygotes carrying missense
mutations the values obtained (without a TMA
challenge) were observed to be in the normal range
25 (data not shown).

 Of the 10 probands with perturbed TMANO:TMA
ratios, DNA was available from 8 individuals; mutation
analysis was performed on these samples. The ethnic
origins as well as the *FM03* genotypes of these are
30 detailed in Table 2: 7 of the probands are of European
Caucasian origin; one is Metis (French Canadian/North
American Indian).

Table 2
FM03 Mutations and Phenotypes
in Eight Probands with TMAuria

Subject (sex)	Age ^a	Ethnic origin	Biochemical phenotype			Genotype ^e	
			TMANO ^c	TMA ^b	TMANO: TMA	Nucleotide change	Amino acid Change
1 ^d (M)	55	Irish/ English	33.90	50.71	39:61	458G>T-(472G)/ 458G>T(472G)	P153L-(E158)/ P153L-(E158)
2 (F)	28	German/ German Irish/ French	17.86	22.73	44:56	154G>A-(472A)- 923A>G/ (472A)-923A>G	A52T-(K158)- E308G/(K158)- E308G
3 (M)	23	German/ Russian	6.05	66.2	8.4:91.6	458G>T-(472G)/ (472G)-1923A>G	P153L-(E158)/ (E158)-E308G- E314X
4 ^c (M)	12	German	11.31	60.24	16:84	458G>T-(472A)/ 458G>T-(472A)	P153L-(K158)/ P153L-(K158)
5 ^c (M)	10	Metis	5.47	65.34	8:92	(472A)-1160G>T/ (472A)-1160G>T	(K158)-R387L/ (K158)-R387L
6 (M)	11	English	6.93	51.90	12:88	458G>T/(472G)/ (472G)-913G>T	P153L-(E158)/ (E158)-E305X
7 (F)	8	French Canadian	12.20	75.8	14:86	458G>T/1474C>T/ (472G/A)	P153LR492W;
8 (M)	2	English	8.36	52.96	14:86	458G>T(472G)/ 458G>T(472G)	(158 E/K)P153L (E158)

^a Age at diagnosis.

^b Units are mmol TMA or TMANO/mol creatinine.

^c Indicates the presence of migraine.

^d Ischemic heart disease.

^e In addition to the mutations, the genotype at polymorphic codon 158 is given in parentheses.

^f Normal values: TMANO > 98% total, TMA < 2% total.

Of previously described mutations, P153L was found to account for nine alleles in this study, while the E305X and R492W mutations were each observed uniquely. M66I, previously observed in the Australian cohort, was not detected in this study. Four new disease-causing alleles were detected: A52T (c.154G→A), E308G (c.923A→G), E314X (c.940G→T), and R387L (c.1160G→T). The substitutions A52T, E314X, and

R387L were not detected on screening 60 control chromosomes from French Canadians. The E308G allele was found to segregate in two probands, homozygous in one compounded with another missense allele (A52T).
5 The G308 allele was detected in 36/198 normal control chromosomes from a Quebec francophone population ($q=0.18$) and in 18 of 118 chromosomes of a Quebec anglophone population ($q=0.15$). We observed E308G in the homozygous state in one French Canadian control.
10 The allele frequency of the G308 allele in probands (0.19) is not statistically different from the background frequency.

Thus, 10 cases of severe TMAuria were confirmed in 28 North Americans with the presenting symptom of
15 malodor. The results are similar to those of the group of George Preti in Philadelphia, who reported that one-third of their referrals for the same complaints were due to TMAuria. While our report defines cases of severe TMAuria, it does not address the possible
20 phenotype of heterozygotes or very mild cases which may be detected by TMA or choline challenge testing. All of the probands studied have TMA oxidation of less than 50% of normal.

A relationship between ablation of FM03
25 activity and symptoms suggestive of disordered biogenic amine metabolism such as hypertension has been noted. In this study, 3 to 10 TMAuria probands also exhibit "labile" hypertension according to their physicians, although one of these individuals (Table
30 2, Subject 1) also has ischemic heart disease which may cause secondary hypertension. Three of the 10 confirmed subjects report the symptoms of classical migraine, a condition noted to be associated with disordered metabolism of biogenic amines such as
35 tyramine. As FM03 is expressed in brain and

metabolizes biogenic amines, this association merits further study.

It is now established that mutation of the *FMO3* gene causes severe TMAuria in North American probands. 100% ascertainment of mutations in cases defined by the admission criteria of the study are reported. Our previous studies of Australian TMAuria probands detected two relatively common mutations (P153L and E305X) in probands of British origin and two rare alleles (M66I, R492W). In this North American cohort, it also found that the mutation P153L is relatively frequent, accounting for 9 of 16 mutant chromosomes. The segregation of the alleles E305X and R492W was also noted. These alleles may have originated in the British Isles.

Four new mutations were detected. The alleles A52T, E314X, and R387L may be "private" mutations, as they have been observed once in the probands of differing ethnic backgrounds. Loss of activity is strongly predicted from truncation of the *FMO3* protein at codon 314 since we know that deletion of even the final 30 amino acids of this 582 residue protein will ablate function *in vitro* (Cashman J (1995) Chem Res Toxicol 8:166-181). While the mutations A52T and R387L have not been expressed, they satisfy several conditions for designation as disease-causing: the changes have not been observed in controls ($n=30$), they are nonconservative substitutions, and no other changes were identified in the probands on sequencing all expressed *FMO3* exons. The A52 and R387 residues appear to be highly conserved within the *FMO* gene family, as seen in Table 3.

Table 3
Amino Acid Sequence Surrounding Human FMO3
Codons 52 and 387; Identity and Variation
among Other FMO Proteins

HFMO3a partial amino acid sequence											Reference
Enzyme	A	E	E	G	R	A52	S	I	Y	K	
RbFMO3	-	Q	(35)
MFMO3	I	(36)
GpFMO5	P	(37)
HFMO5	P	(37)
HFMO1	V	L	.	.	(38)
MFMO1	V	L	.	.	D16215 ^b
PFMO1	V	L	.	.	(39)
RfFMO1	V	L	.	N	(40)
GpFMO1	V	.	D	(41)
RhFMO2	V	.	D	Q	U59453 ^b
HFMO4	S	K	D	.	M	T	R	V	.	.	(42)
RbFMO4	S	K	D	.	M	T	R	V	.	W	(35)

HFMO3 ^a partial amino acid sequence											Reference
Enzyme	V	D	L	Q	S	R387	W	A	A	Q	
RbFMO3	T	.	.	.	A	(35)
MFMO3	T	.	.	.	A	(36)
GpFMO5	S	E	.	.	G	.	.	.	V	.	(37)
HFMO5	S	E	.	.	G	.	.	.	T	.	(37)
HFMO1	G	E	T	.	A	.	.	.	V	R	(38)
MFMO1	G	E	T	.	A	.	.	V	V	.	D16215 ^b
PFMO1	G	.	T	.	A	.	.	.	V	R	(39)
RfFMO1	G	E	T	.	A	.	.	V	V	.	(40)
GpFMO1	.	E	.	.	A	.	.	.	T	R	(41)
RhFMO2	A	E	.	.	A	.	.	V	T	R	U59453 ^b
HFMO4	T	E	.	.	A	.	.	V	T	R	(42)
RbFMO4	T	E	.	.	A	.	.	.	T	R	(35)

^a Abbreviations: Gp, guinea pig; H, human; M, mouse; P, pig; Rb, rabbit; Rh, Rhesus monkey; Rt, rat.

^b GenBank Accession Number.

A comparison of amino acid sequences surrounding these residues in 12 published FMO sequences indicates that R387 is conserved in all 12 while the alanine at codon 52 is conserved in 10 of these 12. The variation at codon 52 seen in the 2 published FMO4 sequences may contribute to substrate specificity differences seen between FMO isoforms.

The G variant at codon 308 was observed in two probands with TMAuria. Subject 2 is an E308G homozygote with a single copy of the A52T missense mutation; Subject 3 is a compound heterozygote, with

E308G and E314X on one homologue and P153L *in trans* (Table 2). The G308 allele was originally observed segregating in normal controls, and it exists as a frequent polymorphism in the Quebec population ($q=0.18$). The E308G polymorphism may also mediate other variations of drug and chemical detoxication. The genotype for Subject 2, who has had complete sequencing of all FMO3 coding exons, is T52-K158-G308 on one allele and A52-K-158-G308 on the other. Previous *in vitro* expression studies on the codon 158 polymorphism indicated the K158 form of the protein is a poorer TMA N-oxygenator than the E158 form. The findings demonstrate that the G308 change on the K158 background functions as a TMAuria mutation. It is possible that these two common changes *in cis* further diminish the activity of the enzyme and render it incapable of compensating for the A52T mutant, resulting in a severe TMAuria phenotype.

As shown in Fig. 1, mutations appear to cluster in exon 7, with three of the four identified changes in this exon occurring within a 10 amino acid "hot spot". These changes fall near or in the areas of high homology described by Ziegler. They are in close proximity to the FATGY signature found in all the mammalian enzymes, while the R387L mutation is found in an area highly conserved among FMOs, strongly suggesting that this substitution affects protein function.

Each of the eight genotypically defined probands studied was ascertained in a different region of North America; presumably these regions differ in background carrier frequencies for FMO3 mutations. Since one of the probands is from Quebec, the overall population frequency question was addressed by screening for known mutations (M66I, P153L, E305X,

R492W) in a Quebec control group. Of 320 control chromosomes, only one carrier for the mutation E305X was detected. Although the number surveyed is small, this predicts that severe TMAuria should be rare in Quebec. Milder variations that may manifest with a precursor load may be more common. This condition may "cluster" in specific regions of the United States and Canada due to "founder effect" and "genetic drift", as occurs in the Melbourne region of Australia. In addition to possible evolutionary advantages in detoxication, there are other hypotheses to account for the maintenance of mutant *FMO3* alleles in the population. These must include other possible selective advantages of heterozygosity, perhaps in blood pressure homeostasis or in metabolism of choline, now recognized to be a vital amine involved in many pathways of intermediary metabolism

EXAMPLE II

Population Specific Polymorphisms of the human *FMO3* gene: Significance for Detoxication

Study Population

The human *FMO3* polymorphisms E158K (c.488 G-A) and V257M (c.769 G-A) were previously identified in a cohort of individuals with trimethylaminuria and normal controls from Melbourne, Australia using single stranded conformational polymorphism (SSCP) screening and sequencing of the *FMO3* gene (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998). Following institutional ethics approval, the frequency of these substitutions were determined in 170 normal control individuals from Quebec (i.e., 110 Francophones and 60 Anglophones) and 50 normal controls from Victoria, Australia. Genomic DNA was prepared from lymphocyte preparations using

standard procedures. Amplified DNA fragments obtained by the polymerase chain reaction were subjected to restriction enzymatic digestion, and visualized by ethidium bromide staining after agarose or polyacrylamide gel electrophoresis. Table 1 illustrates the conditions used for each diagnostic assay.

Table 4
Diagnostic PCR methods created to screen for identified human *FMO3* sequence changes among healthy controls

Allele	Oligonucleotide Sequence	SEQ ID NO:	Restriction Enzyme	Normal Restriction Pattern (bp)	Variant Restriction Pattern (bp)
E158K	2055: 5'-TTG TTC CGG ACA TCA TGT GTA GC -3'	3	<i>HinfI</i>	253	217
	Gen2: 5'-TCC CTG CTG TGG AAG CAT TT -3'	4		76	76
V257M	C1: 5'-TTC CAG AAG TGG CTC CTG GG -3'	5	<i>Hsp92II</i>	97	36
	2005:5'-GCT TGA ATC TTG CAT TCA TCT GC -3'	6		60	72
				60	25

Table 5
Genotype frequencies for the human *FMO3* polymorphisms E158K and V257M in Quebec, Canada and Victoria, Australia

	E158K	V257M
QUEBEC FRANCOPHONE POPULATION	N=109	N=108
	EE:38	VV:101
	EK:56	VM:6
	KK:15	MM:1
QUEBEC ANGLOPHONE POPULATION	N=60	n=58
	EE:16	VV:55
	EK:30	VM:3
	KK:14	
VICTORIA	N=39	N=61
	EE:12	VV:50
	EK:20	VM:9
	KK:78	MM:2

Data Analysis

The statistical significance of the frequency of the human *FMO3* E158K and V257M polymorphisms between Canadians (i.e., Anglophones and Francophones) and Australians was calculated. A chi-squared analysis

was used for human *FMO3* E158K to test the hypothesis that the proportion of chromosomes was not significantly different between the three groups of English Canadians, French Canadians and Australians (χ squared, 2df = 2.52, p = 0.28). The Mantel-Haenszel test for small sample size was used for human *FMO3* V257M. There was no significant difference observed between the presence of the variant alleles in the three cohorts (χ squared, 2df = 4.71, p = 0.095).

10 Chemicals

[³H]-Tyramine was obtained from American Radioactivity Company (St. Louis, MO) and [¹⁴C]-TMA was obtained from Sigma Chemical Co., (St. Louis, MO). All chemicals and reagents were purchased from Aldrich Chemical Co. (Milwaukee, WI) in the highest purity commercially available.

Synthesis

10-(5-Aminopentyl)-2-(trifluoromethyl) phenothiazine (5-APT), its hydroxylamine and *cis* and *trans* oximes were synthesized by a modification of the procedures previously described. The tertiary amine 10-(N,N-dimethylaminopentyl)-2-(trifluoromethyl)phenothiazine (5-DPT) and its N-oxide was also synthesized by a procedure similar to the one previously described. The hydroxylamine and *cis* and *trans* oximes of tyramine were synthesized as previously described (Lin, J et al. (1997) Chem. Res. Toxicol. 10: 842-852).

cDNA-expression and Substrate Analysis

30 The human *FMO3* cDNAs were expressed as maltose binding fusion proteins. Site-directed mutagenesis for the human *FMO3* substitutions E158K and V257M were performed as previously described (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998, Cashman JR et al.

Chem Res Toxicol 10:837-841, 1997). For purposes of comparison, a cDNA construct for the truncation variants E305X (previously reported) and 510X were also prepared as the maltose binding protein fusions.

5 **Subcloning Human FMO1 and FMO3 cDNA into the Maltose Binding Protein Fusion Expression System**

Human FMO3 or FMO1 cDNA was inserted into the expression vector pMAL-c2 and PCR amplification was done in a fashion that allowed for fusion of human
10 FMO3 or FMO1 cDNA at the 3'-end of sequences encoding the maltose binding protein (MBP) as previously described (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998). Each cDNA was individually cloned and confirmed by oligonucleotide sequencing of both
15 strands. The creation and cDNA-expression of wild type Glu 158 and the common polymorphic form Lys158 FMO3-MBPs has been previously described (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998, Cashman JR et al. Chem Res Toxicol 10:837-841, 1997). Another
20 polymorphic form of human FMO3 (i.e., Met 257) was created by oligonucleotide-directed mutagenesis and PCR in a similar fashion as described before. The truncation mutations of FMO3-MBP (E305X and 510X) were created by PCR. The template was the pMAL-2c wild type
25 human FMO3 Glu 158 expression plasmid. At the appropriate position, the forward PCR primer contained a BamHI site that was followed by ATG. The reverse primer changed the codon following the site of truncation to an ochre stop codon, that was also part
30 of a Hind III site. The truncated human FMO3 cDNA was synthesized using Taq polymerase under standard conditions. The PCR fragment was gel purified, digested with BamHI and HindIII and inserted with DNA ligase into the pMAL-c2 vector cut with the same
35 restriction enzymes. Each desired truncation product

was transformed into competent JM109 *E. coli* and plated onto LB-Amp plates. DNA isolated from colonies were shown to contain the desired truncation mutation by sequencing of both strands. The truncation expression plasmids introduced into bacterial strain JM109 were purified by affinity chromatography as previously described.

Electrophoresis and Immunoblotting

Overproduction of the affinity-purified human FMO3-MBP fusion proteins and truncation variants was shown by fractionation on 12% sodium dodecyl sulfate polyacrylic gel electrophoresis (SDS-PAGE). Immunoblots were done according to a previously described procedure using an affinity purified rabbit polyclonal antibody that was directed against the wild type human FMO3-MBP fusion protein.

Enzyme Assays

Assay and analysis of human FMO3-MBP fusion protein and analysis of N-oxygenation activity for the variant enzymes was done by a procedure described previously. The N-oxygenation 5-APT was done using an HPLC method essentially identical to the one described previously for a closely related compound. TMA and tyramine N-oxygenation was determined using radiometric assays described below.

For [³H]-tyramine and [¹⁴C]-TMA, incubations were carried out in 13 x 100 screw-cap culture tubes suspended in a 37°C water bath-metabolic shaker apparatus. The reaction mixture consisted of 10-30 µg of human FMO3-MBP or variant, 0.05 M potassium phosphate buffer (pH = 8.4), 0.8 mM diethylenetriaminepentaacetic acid (DETAPAC), 0.5 mM NADP⁺, 0.5 mM glucose-6-phosphate and 1 IU of glucose-6-phosphate dehydrogenase in a total volume of 0.25 ml. The reaction was initiated by the addition of

radiolabelled substrate to an ice-cold previously equilibrated enzyme solution, capped and incubated at 37°C with constant shaking. For TMA, the reaction was stopped by the addition of 0.25 ml cold CH₃CN containing 0.01 ml trichloroacetic acid (TCA). For tyramine, the reaction was stopped by the addition of 0.25 ml cold MeOH. The incubation mixture was thoroughly mixed and centrifuged and an aliquot was applied to the loading zone of a Whatman Diamond LK6DF TLC plate (Clifton, NJ) previously co-spotted with authentic starting material or products of the specified reaction. For TMA, after air drying, the plate was developed in methanol/chloroform/20% TCA (90:10:0.5, v:v). The following bands (R_f) were visualized by treating with iodine and scraped into scintillation vials for counting: TMA N-oxide (0.32) and TMA (0.08). For tyramine, after air drying, the plate was developed in methanol/dichloromethane/formic acid (20:80:0.2, v:v). The following bands (R_f) were visualized by UV-vis and scraped into scintillation vials for counting: *cis* and *trans* phenylacetone oximes (0.84), tyramine hydroxylamine (0.35) and tyramine (0.04). Each vial was counted in a Beckman LS-2000 scintillation counter and the percent product formation was computed from the data for calculation of the kinetic parameters.

FAD determination

A 0.5 ml aliquot of protein was combined with an equivalent amount of cold acetonitrile, mixed and centrifuged at 12,000 x g. The pellet was washed three times with cold acetonitrile. To the protein pellet was added 50 µl of trifluoroacetic acid, mixed thoroughly and heated in a sealed tube at 55°C for at least 40 min. At the end of the reaction 100 µl of cold acetonitrile was added as well as a sufficient

amount of NH_4OH to neutralize the reaction. A portion of the mixture was injected directly onto a Hitachi HPLC system using a Rainin Microsorb MV C-18 reverse phase column (Rainin, Emeryville, CA). An eluent of water (containing 0.05% HClO_4) for the first 20 min followed by CH_3CN (containing 0.05% HClO_4) thereafter was used to separate FAD from other minor impurities by HPLC. FAD eluted during the first phase of the gradient and was detected at 450 nm with a retention time of 9.2 min. The amount of FAD was determined from a comparison of peak heights from a standard curve of FAD.

RESULTS

Genotype Frequencies

The genotype frequencies for the two prevalent human *FMO3* polymorphisms E158K, and V257M in healthy populations from Quebec, Canada and Victoria, Australia are listed in Table 5. On the basis of statistical analysis, the distribution of the two codon 158 polymorphic variants were found to be similar in Quebec Francophones and Anglophones and in the Australian population. Although the numbers were small, the methionine variant involving codon 257 appeared to be at higher frequency in the Australian population compared with the Quebec population.

Substrate N-Oxygenation

Previous studies showed that non-transformed host bacteria or bacteria transformed with *pMal* alone did not contain any detectable human FMO activity when grown in the presence or absence of IPTG. Previously, it was shown that the relative activity of human *FMO3* MBP and some variants showed varying degrees of TMA N-oxygenation activity (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998, Cashman JR et al. Chem Res Toxicol 10:837-841, 1997). To examine this point more

carefully and to quantify differences among the cDNA-expressed alleles, K_m and V_{max} values were obtained from double reciprocal plots of velocity versus substrate concentration. For the substrates examined, the formation of tertiary amine N-oxide, hydroxylamine or oxime metabolite was a linear function of protein concentration and with incubation time for at least 10 min. As shown by the kinetic constants listed in Table 4, wild-type human *FMO3* MBP efficiently N-oxygenated 5-DPT, TMA and tyramine. Activities for human *FMO1* MBP (a fetal hepatic *FMO* isoform that is expressed in adult kidney and intestine but not in adult liver) were evaluated in a similar kinetic fashion. The kinetic constants are listed in Table 6 for comparison. Human *FMO1* MBP N-oxygenated 5-DPT with a very low K_m value. In contrast, TMA N-oxygenation was significant but the concentration of substrate required for half-maximal activity of *FMO1* MBP was 15-fold greater than that of human *FMO3* MBP. The K_m for human *FMO1* MBP N-oxygenation of TMA compared favorably with that of TMA N-oxygenation for pig *FMO 1* (Cashman J (1995) Chem Res Toxicol 8:166-181).

Table 6
Kinetic Constants for Human FMO1 MBP, FMO3 MBP,
and Variants of Human FMO3 MBP^a

5

Enzyme	K_m	$k_{cat}(min^{-1})$	V_{max}/K_m^c	Substrate
Human FMO1 (n=6)	8.0 ± 2.4	4.0 ± 1.6	0.5	5-DPT
Human FMO3				
(WT) Glu158 (n=3)	154.9 ± 31.8	50.7 ± 1.6	0.33	5-DPT
Lys158 (n=3)	187.5 ± 34.1	44.4 ± 17.7	0.23	5-DPT
Met257 (n=3)	82.3 ± 41.7	13.2 ± 4.6	0.16	5-DPT
510X (n=4)	61.6 ± 12.9	0.09 ± 0.09	0.001	5-DPT
Human FMO1 (n=4)	486 ± 34.8	25.9 ± 3.7	0.05	TMA
Human FMO3				
(WT) Glu158 (n=4)	32.3 ± 7.1	189 ± 21	5.85	TMA
Lys158 (n=4)	206 ± 21	105 ± 3.8	0.51	TMA
Met257 (n=4)	1151 ± 107	28.2 ± 5.1	0.02	TMA
510X (n=4)	1034 ± 139	34.9 ± 14.6	0.03	TMA
Human FMO1 (n=4)	ND ^b			Tyramine
Human FMO3 (n=4)				
(WT) Glu158 (n=4)	231 ± 43.4	110 ± 16.1	0.48	Tyramine
Lys158 (n=4)	941 ± 103	106.1 ± 4.6	0.11	Tyramine
Met257 (n=4)	2164 ± 87	41.3 ± 6.8	0.02	Tyramine
510X (n=4)	1384 ± 110	11.4 ± 6.9	0.008	Tyramine

^aproduct determined by HPLC or radiometric assay. ^bNot Detectable, limit of detection; *trans* oxime, 20 pmol/min/mg of protein. ^c V_{max} is given as nmol/min/nmol of protein, nmol of protein determined on the basis of FAD content as described in the Methods.

A prevalent polymorphic form of human FMO3, FMO3 MBP Lys 158, N-oxygenated 5-DPT, TMA and tyramine with higher K_m values and with a lower V_{max} . Another polymorphic variation of FMO3, FMO3 MBP Met 257, showed significant differences in the kinetic parameters for 5-DPT, TMA and tyramine. For 5-DPT, TMA and tyramine N-oxygenation, the V_{max}/K_m ratios for

human FMO1 MBP were 18-, 249- and 25-fold lower, respectively, than the V_{max}/K_m ratios for the wild type human FMO3 MBP enzyme.

Previously, we have examined the truncation mutation human FMO3 MBP 305X (Treacy EP et al. Hum Molec Genet 7: 839-845, 1998). No detectable N-oxygenase activity was observed for 5-DPT, TMA and tyramine. For comparison, a human FMO3 cDNA construct that was truncated at codon 510 was also analyzed for substrate N-oxygenation activity with 5-DPT, TMA and tyramine. The K_m values for FMO3 MBP 510X were similar to the K_m values observed for FMO3 MBP Met 257, but generally, the V_{max} values were significantly lower than the wild type fusion protein (Table 6).

The Glu 158 wild type human FMO3 MBP enzyme N-oxygenated 5-APT (i.e., the primary amine analog of 5-DPT) with a rate of 117 nmol/min/nmol of protein. As determined by HPLC, the major product was the hydroxylamine. A minor amount of oxime was formed and the ratio of *cis:trans* oxime was 79:21. Human FMO3 MBP Lys158 N-oxygenated the primary amine 5-APT with a rate of 69.1 nmol/min/nmol of protein. As determined by HPLC, the major product was the hydroxylamine. A minor product was the oxime and it was formed in a ratio of *cis:trans* oxime 81:19. By comparison, the human FMO3 Met 257 enzyme N-oxygenated the primary amine 5-APT with a rate of 28.1 nmol/min/nmol of protein. As determined by HPLC, the major product was the hydroxylamine. A minor product was the oxime and it was formed with a *cis:trans* stereoselectivity of 80:20. Human FMO1 MBP did not significantly N-oxygenate tyramine or 5-APT.

Because 5-APT was present in large excess, the rates were determined at saturating substrate concentration and represent apparent V_{max} values. We

examined a number of human *FMO3* MBP truncation mutations introduced between codon 305 and the wild type length. There was no significant N-oxygenase activity observed for the human *FMO3* cDNA construct with a truncation less than codon 510. Similar experiments have been conducted with pig *FM01* MBP and a similar structure-function relationship has been observed (Cashman J (1995) Chem Res Toxicol 8:166-181).

EXAMPLE III

TMAuria and variation of the *FMO3* Gene Dietary and Pharmacogenomic Implications

The relationships between choline (precursor of trimethylamine (TMA)), TMA and choline products (phosphatidylcholine, acetylcholine) in individuals with trimethylaminuria (TMAuria, OMIM 602079) and normal controls may be characterized to determine the optimum dietary treatment and choline requirements for individuals with TMAuria. Patients with TMAuria may be treated with riboflavin (co-factor). The consequences of polymorphic variants of the *FMO3* gene for TMA metabolism may be understood and applied the metabolism of other nitrogen-containing *FMO3* substrates and common medications.

Treatment of individuals with TMAuria

Individuals with TMAuria may be supplemented with riboflavin. The vitamin riboflavin (vitamin B2) is the cofactor for the FMO FAD-dependent enzyme family. Supplementation with riboflavin may increase the residual activity of *FMO3* and other FMO isoforms known to oxygenate TMA as observed in other disorders. FMO enzymes are known to be inducible by substrate loads and hormones (Cashman J (1995) Chem Res Toxicol 8:166-181). Affected probands may be studied. The

patients may take a once daily dose of riboflavin 200 mg for a period of two months. A baseline urine sample for TMA and TMA N-oxide (FAB-MS) may be measured and at intervals of 4, 6 and 8 weeks for measurement of effect.

The three *FMO3* polymorphisms (E158K, V257M and E308G) may be characterized further. These polymorphisms may exert a clinical effect on TMA oxygenation, applicable, perhaps to other *FMO3* substrates including commonly used medication. DNA samples from 70 extended French Canadian families (450 individuals) from the Chicoutimi region of Quebec; (Hypertension sib-pair analysis study). have been genotyped for the three polymorphisms, and approximately 20% of individuals are homozygous or compound heterozygous for one or more of these polymorphisms. Individuals with these genotypes may exhibit a difference in TMA oxidation (used as a marker of decreased N-oxygenation). A rapid FAB-MS assay for TMA and its N-oxide may be used to assess TMA and TMANO in the informative cases. A detailed drug history may also be taken from these individuals to assess whether they are taking other medications likely to be *FMO3* substrates for future pharmacogenetic studies.

From a large cohort of individuals with TMAuria worldwide (confirmed cases in North America and Australians) plasma choline, acetylcholine, phosphatidylcholine, and TMA and TMANO may be measured in urine in controls (adult and pediatric). This may also be performed for individuals with untreated TMAuria and individuals on a low choline diet. Dietary evaluation of daily choline intake (free choline, lecithin, phosphocholine, phosphatidylcholine) may be analyzed using a 3-day dietary recall for the days

prior to the blood test. The choline content of foods may be analyzed and correlated with TMA oxidation results and genotype. As choline is oxidized to betaine, important in homocysteine remethylation, total homocysteine may also be measured in individuals on a choline-restricted diet. TMA and its N-oxide may be measured using the stable isotope FAB method using a 5 cc urine aliquot. ¹⁵N TMA and TMANO may be used as internal standards. Choline, phosphatidylcholine and acetylcholine may be measured from a blood sample extract using electrospray tandem mass spectrometry. Measurements of urinary TMA and its N-oxide may be used also to determine the riboflavin effect and the in vivo effects of polymorphisms, as mentioned above.

15 Statistical analysis

From data derived from frequency of genotypes as illustrated in Table 7, the number of individuals that will have informative genotypes may be estimated (presence of greater than 2 polymorphisms (cis or trans)); 2 variant alleles: (30%), 3: (14.3%), 4: (2.7%), 5 or more: (less than 1%). The estimated sample size of 76 (allowing for non-compliance) may permit adequate comparison between cases and controls. The Student's t test may be used to test the hypothesis that the group of individuals with variant genotypes have a significantly lower mean percentage of N-oxidation than controls. If the sample size permits, the study group may be divided into those that carry 2, 3 or 4 polymorphic alleles and the t-test performed on the distribution of N-oxygenation for each of these categories compared to the central distribution.

These studies may improve our understanding and treatment of TMAuria and our understanding of the

WO 01/23603

PCT/CA00/01118

- 36 -

medical relevance of the polymorphisms for TMA metabolism applicable to other FMO substrates.

Table 7
Genotypes of Possible Pharmacogenetic Significance
Observed in Chicoutimi Region

Polymorphisms	Allele 1	Allele 2
E158K	2	2
V257M	1	1
E308G	1	1
E158K	2	2
V257M	1	1
E308G	1	2
E158K	1	2
V257M	1	1
E308G	2	2
E158K	2	2
V257M	1	1
E308G	2	2
E158K	1	2
V257M	1	2
E308G	1	2
E158K	1	2
V257M	2	1
E308G	1	2
E158K	1	1
V257M	2	2
E308G	1	1
E158K	1	2
V257M	2	1
E308G	1	1
E158K	1	1
V257M	2	2
E308G	1	2
E158K	2	2
V257M	1	2
E308G	1	1
E158G	2	2
V257M	1	2
E308G	2	2
E158G	1	2
V257M	1	2
E308G	1	2

5 1 designates wild allele, 2 designates variant allele

EXAMPLE IV

**Polymorphisms of the flavin containing monooxygenase 3
(FMO3) gene as a risk factor for essential
hypertension in French Canadian**

5 Interindividual variation in blood pressure is
genetically determined resulting from the interaction
of susceptibility alleles and environmental exposures.
We previously documented that the rare mutations of
the *FMO3* gene (a phase I xenobiotic detoxicating gene)
10 cause the inborn error of metabolism Trimethylaminuria
and may predispose affected individuals to essential
hypertension (HBP) and abnormal catecholamine
polymorphisms (E158L (c.488 G-A), E308G (c.923 A-G)
and V257M (c. 769 G-A)) and haplotypes were phased in
15 70 extended families from Quebec. Of 8 possible
haplotypes, 6 were observed. These *in vivo* and *in*
vitro data indicate that these polymorphisms in
particular haplotypes confer functional
pharmacogenetic effects.

20 In accordance with the present invention, now
is reported a case control study of the association
between variant *FMO3* haplotypes and early onset HBP (<
55 years) in a Quebec French Canadian population. The
distribution of haplotypes was determined between (1)
25 HBP probands ($n = 60$), (2) HBP relatives ($n = 229$),
(3) non-HBP control relatives ($n = 145$) and (4)
unrelated non-HBP French Canadian controls ($n = 133$).
There was a significant difference noted between the
distribution of haplotypes between groups (1) and (4)
30 ($p = 0.026$). The presence of one variant haplotype in
individuals conferred a relative risk of HBP of 1.22
variant haplotypes. These data suggest that the above
common polymorphisms of the *FMO3* gene in the presence
of adverse environmental exposures may increase the
35 risk of individuals in Quebec to early onset HBP.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to
5 cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention
10 pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A method for detecting an altered metabolism of a substrate of a flavin-containing monooxygenase (FMO) enzyme or an isoform thereof in a patient, the method comprising detecting at least one of a mutation and a polymorphic variant of a gene encoding said FMO enzyme in a sample from said patient, whereby said at least one of said mutation and said polymorphic variant is indicative of an altered metabolism for said substrate.
2. A method for detecting a susceptibility of a patient to a substrate of a flavin-containing monooxygenase (FMO) enzyme or an isoform thereof in a patient, the method comprising detecting at least one of a mutation and a polymorphic variant of a gene encoding said FMO enzyme in a sample from said patient, whereby said at least one of said mutation and said polymorphic variant is indicative of a susceptibility to said substrate.
3. A method for detecting a predisposition of a patient to a disorder associated with an (adverse) exposure to a heteroatom-containing chemical compound, an intermediate or a metabolite thereof associated with carcinogenesis or having a toxic, pro-carcinogenic or carcinogenic potential, said method comprising detecting at least one of a polymorphic variant and a mutation of a gene encoding a flavin-containing monooxygenase (FMO) enzyme or an isoform thereof in a sample from said patient, whereby said at least one said polymorphic variant and said mutation

is indicative of exposure to the chemical compound, the intermediate or the metabolite thereof.

4. A method for detecting a predisposition to hypertension in a patient, said method comprising detecting at least one of a mutation and a polymorphic variant of a gene encoding said FMO enzyme in a sample from said patient, whereby said at least one of said mutation and said polymorphic variant is indicative of a predisposition to hypertension.

5. A method according to anyone of claims 1 to 4, wherein said at least one of said mutation and said polymorphic variant inactivates partially or totally the activity of said FMO enzyme.

6. A method according to claim 5, wherein said FMO enzyme consists of isoform 3 (FMO3).

7. A method according to claim 6, wherein the polymorphic variant comprises a polymorphic variant from the group consisting of E158K, V257M and E308G, and the mutation comprises a mutation from the group consisting of P153L, E305X, M66I, E314X, R492W, A52T and R387L.

8. A method according to claim 7, wherein the altered metabolism is associated with an idiosyncratic reaction to the substrate.

9. A method according to claim 8, wherein the altered metabolism is associated with a disorder.

10. A method according to claim 9, wherein the disorder is a cancer.

11. A method according to claim 7, wherein the substrate is a xenobiotic or an endogenous material relative to said individual.

12. A method according to claim 11, wherein said xenobiotic is a drug, a food additive, a pesticide, a plant toxin, an organic chemical compound or an aromatic amine.

13. A method according to claim 12, wherein the substrate is a biogenic amine contained in diet of said individual.

14. A method according to claim 13, wherein said biogenic amine is a tertiary amine.

15. A method according to claim 14, wherein said tertiary amine is trimethylamine (TMA), tyramine or catecholamine.

16. A method according to claim 14, wherein the tertiary amine is trimethylamine (TMA) and wherein the disorder is trimethylaminuria (TMAuria).

17. A method for the treatment of an individual having a disorder associated with an altered activity of a flavin-containing monooxygenase FMO enzyme or an isoform thereof, the method comprising supplementing the individual with riboflavin to increase the altered activity of the FMO enzyme or isoform thereof.

18. A method according to claim 17, wherein the isoform consists of isoform 3 (FMO3).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number
WO 01/23603 A3(51) International Patent Classification⁷: C12Q 1/68, A61P 43/00, A61K 31/525

(21) International Application Number: PCT/CA00/01118

(22) International Filing Date:
28 September 2000 (28.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/156,934 30 September 1999 (30.09.1999) US

(71) Applicant (for all designated States except US): MCGILL UNIVERSITY [CA/CA]; 845 Sherbrooke West, Montréal, Québec H3A 2T5 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TREACY, Eileen, P. [CA/CA]; 462 Elm Avenue, Westmount, Québec H3Y 3J1 (CA). AKERMAN, Beverly, R. [CA/CA]; 6076 Notre-Dame de Grace, Montréal, Québec H4A 1N2 (CA).

(74) Agents: COTE, France et al.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College Avenue, Montréal, Québec H3A 2Y3 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments


(88) Date of publication of the international search report:
4 April 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HUMAN FMO3 GENE MUTATIONS AND POLYMORPHISMS, AND USES THEREOF

(57) Abstract: The present invention relates to human FMO3 gene polymorphisms and more particularly to uses thereof in the diagnosis of trimethylaminuria. There is provided a method for detecting an altered metabolism of a substrate of a flavin-containing monooxygenase form 3 (FMO3) enzyme in an individual, detecting a susceptibility of an individual to a substrate of the FMO3 enzyme in an individual, and for detecting a predisposition of an individual to a disorder associated with an exposure to a heteroatom-containing chemical compound, by detecting mutations or polymorphic variants of the FMO3 gene.

WO 01/23603 A3



1596-nt (532 amino acids)

Fig. 1

*CpG site

DECLARATION AND POWER OF ATTORNEY FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63)		Attorney Docket No.:	SWA4338P0080US
		First Named Inventor:	Eileen P. Treacy et al.
		<i>COMPLETE IF KNOWN</i>	
		Application Number:	10/089,491
		Int'l Appln. No.:	PCT/CA00/01118
		Filing Date:	March 28, 2002
<input type="checkbox"/> Declaration Submitted With Initial Filing		Int'l Filing Date:	September 28, 2000
		Group Art Unit:	Unassigned
		Examiner Name:	Not Yet Known
<input checked="" type="checkbox"/> Declaration Submitted After Initial Filing (surcharge (37 CFR 1.16(a)) required)			

As a below-named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first, and sole inventor (if only one name is listed) or an original, first and joint inventor (if plural names are listed) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **Human FMO3 Gene Mutations and Polymorphisms, and Uses Thereof**, the specification of which:

- ☐ is attached hereto; or
- ☒ was filed on March 28, 2002 as Application Serial No. 10/089,491 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information which is material to patentability as defined in 37 CFR. 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Numbers	Country	Foreign Filing Date (MM/DD/YY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
CA00/01118	PCT	09/28/00	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- ☐ Additional foreign application numbers are listed on a supplemental priority data sheet attached hereto.

I hereby claim the benefit of any United States application(s) listed below.

Application Number(s)	Filing Date	<input type="checkbox"/> Additional application numbers are listed on a supplemental priority data sheet attached hereto.
60/156,934	September 30, 1999	

The undersigned hereby authorizes the U.S. attorney(s) or agent(s) named herein to accept and follow instructions from the assignee, if any, of the undersigned or from _____ as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney(s) or agent(s) and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney(s) or agent(s) named herein will be so notified by the undersigned.

As a named inventor, I hereby appoint the following registered practitioner(s) identified by Customer No. 32116 to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, whose firm name, mailing address, telephone number, and facsimile number for this application are:

Wood, Phillips, Katz, Clark & Mortimer
Citicorp Center, Suite 3800
500 West Madison Street
Chicago, Illinois 60661-2511

Telephone (312) 876-1800
Facsimile (312) 876-2020

Customer Number (32116)
and/or Bar Code Label:

--

I hereby declare that all statements made herein of my knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: <u>Eileen P. Treacy</u>	
Citizenship: <u>Canada</u>	
Residence: <u>21 Sydenham Court, Sydenham Rd, Bellsbridge, Dublin 4 PQC</u>	
Post Office Address (if different):	
Signature: <u>Eileen Treacy</u>	Date: <u>28/8/02</u>
<input type="checkbox"/> A petition has been filed for this unsigned inventor.	

201

Name of Additional Inventor, if any: Beverly R. Akerman	
Citizenship: Canada	
Residence: 3466 Belmore Avenue, Montreal, Quebec H4B 2B9 CA	
Post Office Address (if different):	
Signature: <i>B Akerman</i>	Date: <i>June 14, 2002</i>
<input type="checkbox"/> A petition has been filed for this unsigned inventor.	

Name of Additional Inventor, if any:	
Citizenship:	
Residence:	
Post Office Address (if different):	
Signature:	Date:
<input type="checkbox"/> A petition has been filed for this unsigned inventor.	

Name of Additional Inventor, if any:	
Citizenship:	
Residence:	
Post Office Address (if different):	
Signature:	Date:
<input type="checkbox"/> A petition has been filed for this unsigned inventor.	

Name of Additional Inventor, if any:	
Citizenship:	
Residence:	
Post Office Address (if different):	
Signature:	Date:
<input type="checkbox"/> A petition has been filed for this unsigned inventor.	

SEQUENCE LISTING

<110> TREACY, Eileen P.
AKERMAN, Beverly R.
McGill University

<120> HUMAN FMO3 GENE MUTATIONS AND
POLYMORPHISMS, AND USES THEREOF

<130> 1770-235PCT

<150> US 60/156,934

<151> 1999-09-30

<160> 10

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 23

<212> DNA

<213> artifical sequence

<220>

<223> sequence to be used as a primer

<400> 1

gacctgatca gtatactcat tta

23

<210> 2

<211> 23

<212> DNA

<213> artifical sequence

<220>

<223> sequence to be used as a primer

<400> 2

aatgggaagt ctgggaaaca ctt

23

<210> 3

<211> 23

<212> DNA

<213> artifical sequence

<220>

<223> sequence to be used as a primer

<400> 3

ttgttcgga catcatgtgt agc

23

<210> 4

<211> 20

<212> DNA

<213> artifical sequence

<220>

<223> sequence to be used as a primer

<400> 4
tccctgctgt ggaagcattt 20

<210> 5
<211> 20
<212> DNA
<213> artifical sequence

<220>
<223> sequence to be used as a primer

<400> 5
ttccagaagt ggctcctggg 20

<210> 6
<211> 23
<212> DNA
<213> artifical sequence

<220>
<223> sequence to be used as a primer

<400> 6
gcttgaatct tgcattcatc tgc 23

<210> 7
<211> 23
<212> DNA
<213> artifical sequence

<220>
<223> sequence to be used as a primer

<400> 7
ccttatcaat ttatatatgg acc 23

<210> 8
<211> 23
<212> DNA
<213> artifical sequence

<220>
<223> sequence to be used as a primer

<400> 8
ggaccttgta actaggatta ttg 23

<210> 9
<211> 23
<212> DNA
<213> artifical sequence

<220>
<223> sequence to be used as a primer

<400> 9
gaaatgccat actgaccca tgg 23

<210> 10
<211> 23
<212> DNA